

- shadows in the visual behaviour of tsetse flies. *Proc. Biol. Sci.* 271(Suppl 3): S16-S17. <https://doi.org/10.1098/rsbl.2003.0121>
- [10] Kristjanson PM, Swallow BM, Rowlands GJ, Kruska RL, de Leeuw PN. (1999). Measuring the costs of African animal Trypanosomiasis, the potential benefits of control and returns to research. *Agric. Sys.* 59: 79-98. [https://doi.org/10.1016/S0308-521X\(98\)00086-9](https://doi.org/10.1016/S0308-521X(98)00086-9)
- [11] Schofield CJ, Kabayo JP. (2008). Trypanosomiasis vector control in Africa and Latin America. *Parasites & Vectors* 1: 24. <https://doi.org/10.1186/1756-3305-1-24>
- [12] Muturi CN, Ouma JO, Malele II, Ngure RM, Rutto JJ, Mithöfer KM, Enyaru J, Masiga DK. (2011). Tracking the feeding patterns of tsetse flies (*Glossina* genus) by Masiga, analysis of bloodmeals using mitochondrial cytochromes genes. *PLoS One* 6(2): e17284. <https://doi.org/10.1371/journal.pone.0017284>
- [13] Ruiz-Postigo JA, Franco JR, Lado M, Simarro PP. (2012). Human African trypanosomiasis in South Sudan: How can we prevent a new epidemic? *PLoS Negl. Trop. Dis.* 6: e1541. <https://doi.org/10.1371/journal.pntd.0001541>
- [14] Funk S, Nishiura H, Heesterbeek H, Edmunds WJ, Checchi F. (2013). Identifying Transmission cycles at the human-animal interface: The role of animal reservoirs in maintaining Gambiense Human African Trypanosomiasis. *PLoS. Comput. Biol.* 9. <https://doi.org/10.1371/journal.pcbi.1002855>
- [15] Lillesand T, Kiefer WR, Chipman J. (2015). Remote sensing and image interpretation. 7th Edition, John Wiley and Sons, New York.
- [16] Burough P. (2001). Principles of geographical information system, spatial information systems and Geostatistics. New York: Oxford University Press, p. 20.
- [17] Elias S, Tim-Robinson ND. (2007). GIS and multiple-criteria evaluation for the optimisation of tsetse fly eradication programmes. *Environmental Monitoring and Assessment* 124(1-3): 89-103. <https://doi.org/10.1007/s10661-006-9210-0>
- [18] Mugenyi AW. (2014). The spatial distribution of tsetse (Diptera: Glossinidae) within the *Trypanosoma brucei rhodesiense* focus of Uganda. Phd Thesis, University of Edinburgh.
- [19] Palaniyandi, M. (2012). The role of remote sensing and GIS for spatial prediction of vector-borne diseases transmission: A systematic review. *Journal of Vector Borne Diseases* 49(4): 197-204.
- [20] Cecchi G, Mattioli RC, Slingenbergh J, Rocque SD. (2008). Land cover and tsetse fly distributions in sub-Saharan Africa. *Medical and Veterinary Entomology* 22(4): 364-373. <https://doi.org/10.1111/j.1365-2915.2008.00747.x>
- [21] Pagabeleguem S, Ravel S, Dicko AH, Vreysen MJB, Parker A, Takac P. (2016). Influence of temperature and relative humidity on survival and fecundity of three tsetse strains. *Parasites and Vectors* 9: 520. <https://doi.org/10.1186/s13071-016-1805-x>
- [22] Bouyer J, Dicko AH, Cecchi G, Ravel S, Guerrini L, Solano P. (2015). Mapping landscape friction to locate isolated tsetse populations that are candidates for elimination. *Proc Natl Acad Sci.* 112(47): 14575-14580. <https://doi.org/10.1073/pnas.1516778112>
- [23] Tirmizi ST, Tirmizi SRI. (2017). GIS based risk assessment of oil and gas infrastructure in Sindh, Pakistan. *Environmental and Earth Sciences Research Journal* 4(3): 55-59. <https://doi.org/10.18280/eesrj.040301>
- [24] Ngonyoka A, Gwakisa PS, Estes AB, Nnko HJ, Hudson PJ, Cattadori IM. (2017). Variation of tsetse fly abundance in relation to habitat and host presence in the Maasai steppe, Tanzania. *Journal of Vector Ecology* 42(1): 34-43. <https://doi.org/10.1111/jvec.12237>
- [25] Alderton S, Macleod ET, Anderson NE, Palmer G, Machila N, Simuunza M, Welburn SC, Atkinson PM. (2018). An agent-based model of tsetse fly response to seasonal climatic drivers: Assessing the impact on sleeping sickness transmission rates. *PLoS Negl Trop Dis* 2: 10. <https://doi.org/10.1371/journal.pntd.0006188>
- [26] Onyekwelu KC, Ejezie FE, Eze AA, Ikekpeazu JE, Ezech RC, Edeh GC. (2017). Prevalence of trypanosome infection in tsetse flies from Oji river and emene axis of Enugu State, Nigeria: A preliminary report. *Trop Parasitol* 7(2): 98-102. https://doi.org/10.4103/tp.TP_14_17
- [27] Moore N, Messina J. (2010). A landscape and climate data logistic model of tsetse distribution in Kenya. *Plos ONE* 5(7): e11809. <https://doi.org/10.1371/journal.pone.0011809>
- [28] National Population Commission (2009). Federal Republic of Nigeria Official Gazette. The Federal Government Printers Lagos, Nigeria, pp. 493-500.
- [29] Mather PM. (1999). Computer Processing of Remotely Sensed Images – An Introduction. John Wiley and sons, Wiley, Toronto.
- [30] Syed A, Abdulla SMS. (2002). Assessing desert vegetation cover using remotely sensed data: A case study from the state of Qatar. www.gisdevelopment.net.
- [31] Anderson TA. (2002). Planning the built environment. Planners Press American Planners Association. Chicago, Washinton D.C., pp. 1-214.